

A note on interval edge-colorings of graphs

R.R. Kamalian^{ab*}, P.A. Petrosyan^{ac†}

^aInstitute for Informatics and Automation Problems,
National Academy of Sciences, 0014, Armenia

^bDepartment of Applied Mathematics and Informatics,
Russian-Armenian State University, 0051, Armenia

^cDepartment of Informatics and Applied Mathematics,
Yerevan State University, 0025, Armenia

An edge-coloring of a graph G with colors $1, 2, \dots, t$ is called an interval t -coloring if for each $i \in \{1, 2, \dots, t\}$ there is at least one edge of G colored by i , and the colors of edges incident to any vertex of G are distinct and form an interval of integers. In this paper we prove that if a connected graph G with n vertices admits an interval t -coloring, then $t \leq 2n - 3$. We also show that if G is a connected r -regular graph with n vertices has an interval t -coloring and $n \geq 2r + 2$, then this upper bound can be improved to $2n - 5$.

Keywords: edge-coloring, interval coloring, bipartite graph, regular graph

1. Introduction

All graphs considered in this paper are finite, undirected, and have no loops or multiple edges. Let $V(G)$ and $E(G)$ denote the sets of vertices and edges of G , respectively. An (a, b) -biregular bipartite graph G is a bipartite graph G with the vertices in one part all having degree a and the vertices in the other part all having degree b . A partial edge-coloring of G is a coloring of some of the edges of G such that no two adjacent edges receive the same color. If α is a partial edge-coloring of G and $v \in V(G)$, then $S(v, \alpha)$ denotes the set of colors of colored edges incident to v .

An edge-coloring of a graph G with colors $1, 2, \dots, t$ is called an interval t -coloring if for each $i \in \{1, 2, \dots, t\}$ there is at least one edge of G colored by i , and the colors of edges incident to any vertex of G are distinct and form an interval of integers. A graph G is interval colorable, if there is $t \geq 1$ for which G has an interval t -coloring. The set of all interval colorable graphs is denoted by \mathfrak{N} . For a graph $G \in \mathfrak{N}$, the greatest value of t for which G has an interval t -coloring is denoted by $W(G)$.

*email: rrkamalian@yahoo.com.

†email: pet_petros@{ipia.sci.am, ysu.am, yahoo.com}

The concept of interval edge-coloring was introduced by Asratian and Kamalian [2]. In [2, 3] they proved the following theorem.

Theorem 1 *If G is a connected triangle-free graph and $G \in \mathfrak{N}$, then*

$$W(G) \leq |V(G)| - 1.$$

In particular, from this result it follows that if G is a connected bipartite graph and $G \in \mathfrak{N}$, then $W(G) \leq |V(G)| - 1$. It is worth noting that for some families of bipartite graphs this upper bound can be improved. For example, in [1] Asratian and Casselgren proved the following

Theorem 2 *If G is a connected (a, b) -biregular bipartite graph with $|V(G)| \geq 2(a + b)$ and $G \in \mathfrak{N}$, then*

$$W(G) \leq |V(G)| - 3.$$

For general graphs, Kamalian proved the following

Theorem 3 [6]. *If G is a connected graph and $G \in \mathfrak{N}$, then*

$$W(G) \leq 2|V(G)| - 3.$$

The upper bound of Theorem 3 was improved in [5].

Theorem 4 [5]. *If G is a connected graph with $|V(G)| \geq 3$ and $G \in \mathfrak{N}$, then*

$$W(G) \leq 2|V(G)| - 4.$$

On the other hand, in [7] Petrosyan proved the following theorem.

Theorem 5 *For any $\varepsilon > 0$, there is a graph G such that $G \in \mathfrak{N}$ and*

$$W(G) \geq (2 - \varepsilon)|V(G)|.$$

For planar graphs, the upper bound of Theorem 3 was improved in [4].

Theorem 6 [4]. *If G is a connected planar graph and $G \in \mathfrak{N}$, then*

$$W(G) \leq \frac{11}{6}|V(G)|.$$

In this note we give a short proof of Theorem 3 based on Theorem 1. We also derive a new upper bound for the greatest possible number of colors in interval edge-colorings of regular graphs.

2. Main results

Proof of Theorem 3. Let $V(G) = \{v_1, v_2, \dots, v_n\}$ and α be an interval $W(G)$ -coloring of the graph G . Define an auxiliary graph H as follows:

$$V(H) = U \cup W, \text{ where}$$

$$U = \{u_1, u_2, \dots, u_n\}, W = \{w_1, w_2, \dots, w_n\} \text{ and}$$

$$E(H) = \{u_i w_j, u_j w_i \mid v_i v_j \in E(G), 1 \leq i \leq n, 1 \leq j \leq n\} \cup \{u_i w_i \mid 1 \leq i \leq n\}.$$

Clearly, H is a connected bipartite graph with $|V(H)| = 2|V(G)|$.

Define an edge-coloring β of the graph H in the following way:

- (1) $\beta(u_i w_j) = \beta(u_j w_i) = \alpha(v_i v_j) + 1$ for every edge $v_i v_j \in E(G)$,
- (2) $\beta(u_i w_i) = \max S(v_i, \alpha) + 2$ for $i = 1, 2, \dots, n$.

It is easy to see that β is an edge-coloring of the graph H with colors $2, 3, \dots, W(G) + 2$ and $\min S(u_i, \beta) = \min S(w_i, \beta)$ for $i = 1, 2, \dots, n$. Now we present an interval $(W(G) + 2)$ -coloring of the graph H . For that we take one edge $u_{i_0} w_{i_0}$ with $\min S(u_{i_0}, \beta) = \min S(w_{i_0}, \beta) = 2$, and recolor it with color 1. Clearly, such a coloring is an interval $(W(G) + 2)$ -coloring of the graph H . Since H is a connected bipartite graph and $H \in \mathfrak{N}$, by Theorem 1, we have

$$W(G) + 2 \leq |V(H)| - 1 = 2|V(G)| - 1, \text{ thus}$$

$$W(G) \leq 2|V(G)| - 3.$$

□

Theorem 7 *If G is a connected r -regular graph with $|V(G)| \geq 2r + 2$ and $G \in \mathfrak{N}$, then*

$$W(G) \leq 2|V(G)| - 5.$$

Proof. In a similar way as in the prove of Theorem 3, we can construct an auxiliary graph H and to show that this graph has an interval $(W(G) + 2)$ -coloring. Next, since H is a connected $(r + 1)$ -regular bipartite graph with $|V(H)| \geq 2(2r + 2)$ and $H \in \mathfrak{N}$, by Theorem 2, we have

$$W(G) + 2 \leq |V(H)| - 3 = 2|V(G)| - 3, \text{ thus}$$

$$W(G) \leq 2|V(G)| - 5.$$

□

References

- [1] A.S. Asratian, C.J. Casselgren, On interval edge colorings of (α, β) -biregular bipartite graphs, *Discrete Mathematics* 307 (2006) 1951-1956.
- [2] A.S. Asratian, R.R. Kamalian, Interval colorings of edges of a multigraph (in Russian), *Appl. Math.* 5 (1987) 25-34.
- [3] A.S. Asratian, R.R. Kamalian, Investigation on interval edge-colorings of graphs, *J. Combin. Theory Ser. B* 62 (1994) 34-43.
- [4] M.A. Axenovich, On interval colorings of planar graphs, *Congr. Numer.* 159 (2002) 77-94.
- [5] K. Giaro, M. Kubale, M. Malafiejski, Consecutive colorings of the edges of general graphs, *Discrete Math.* 236 (2001) 131-143.
- [6] R.R. Kamalian, Interval edge-colorings of graphs, *Doctoral Thesis*, Novosibirsk, 1990.
- [7] P.A. Petrosyan, Interval edge-colorings of complete graphs and n -dimensional cubes, *Discrete Mathematics* 310 (2010) 1580-1587.